## IN THE CLAIMS

- 1. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising the steps of
  - i) contacting a test compound with a PPARA polypeptide, and
  - ii) detecting detect binding of said test compound to said PPARA polypeptide.
- 2. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising the steps of
  - i) determining the activity of a PPARA polypeptide at a certain concentration of a test compound or in the absence of said test compound, and
  - ii) determining the activity of said polypeptide at a different concentration of said test compound.
- 3. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising the steps of
  - i) determining the activity of a PPARA polypeptide at a certain concentration of a test compound, and

- ii) determining the activity of a PPARA polypeptide at the presence of a compound known to be a regulator of a PPARA polypeptide.
- 4. (currently amended) The method of claim 1 any of claims 1 to 3, wherein the step of contacting is in or at the surface of a cell.
- 5. (currently amended) The method of claim 1 any-of-claims 1 to-3, wherein the cell is in vitro.
- 6. (currently amended) The method of <u>claim 1</u> <del>any of claims 1 to 3</del>, wherein the step of contacting is in a cell-free system.
- 7. (currently amended) The method of <u>claim 1</u> <del>any of claims 1 to 3</del>, wherein the polypeptide is coupled to a detectable label.
- 8. (currently amended) The method of <u>claim 1</u> <del>any of claims 1 to 3</del>, wherein the compound is coupled to a detectable label.
- 9. (currently amended) The method of <u>claim 1</u> any of claims 1 to 3, wherein the test compound displaces a ligand which is first bound to the polypeptide.
- 10. (currently amended) The method of claim 1 any of claims 1 to 3, wherein the polypeptide is attached to a solid support.
- 11. (currently amended) The method of claim 1 any of claims 1 to 3, wherein the compound is attached to a solid support.
- 12. (currently amended) A method of screening for therapeutic agents useful in the treatment of a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising the steps of

- i) contacting a test compound with a PPARA polynucleotide, and
- ii) <u>detecting</u> <del>detect</del> binding of said test compound to said PPARA polynucleotide.
- 13. (original) The method of claim 12 wherein the nucleic acid molecule is RNA.
- 14. (original) The method of claim 12 wherein the contacting step is in or at the surface of a cell.
- 15. (original) The method of claim 12 wherein the contacting step is in a cell-free system.
- 16. (original) The method of claim 12 wherein polynucleotide is coupled to a detectable label.
- 17. (original) The method of claim 12 wherein the test compound is coupled to a detectable label.
- 18. (currently amended) A method of diagnosing a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal comprising the steps of
- i) determining the amount of a PPARA polynucleotide in a sample taken from said mammal, and
- ii) determining the amount of PPARA polynucleotide in healthy and/or diseased mammals.

## 19-20. (canceled)

21. (currently amended) A pharmaceutical composition for the treatment of a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases,

gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising a therapeutic agent which regulates the activity of a PPARA polypeptide, wherein said therapeutic agent is

- i) a small molecule,
- ii) an RNA molecule,
- iii) an antisense oligonucleotide,
- iv) a polypeptide,
- v) an antibody, or
- vi) a ribozyme.
- 22. (currently amended) A pharmaceutical composition for the treatment of a disease selected from the group comprised in a group of diseases consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising a PPARA polynucleotide.
- 23. (currently amended) A pharmaceutical composition for the treatment of a disease selected from the group comprised in a group of diseases—consisting of cardiovascular diseases, gastroenterological diseases, cancer, inflammation, hematological diseases, respiratory diseases, neurological diseases and urological diseases in a mammal, comprising a PPARA polypeptide.

24-26. (canceled)